

ESCRS 2024

SPEAKERS PROGRAM

You don't want to miss this!

Join our **Expert Panel** and learn about iStent *inject*[®] W, a **PROVEN SAFE** technology for an early intervention preserving Patients' Functional Vision¹ and Quality of Life²



GLAUKOS BOOTH
D56, Hall 7

SATURDAY SEPT. 7th

- 11:00 **Safety First:
Optimizing Surgical Order
for Glaucoma Treatment**
Florian Rüfer
- 12:30 **The Test of Time:
Long-term Data Proves
Safety and Efficacy**
Fritz Hengerer
- 14:00 **Reducing surgical Risks:
The Impact of iStent *inject*[®]
on Trabeculectomies**
Laura Morales
- 15:00 **Proven Over Time:
iStent[®]'s Long-Term Safety
and Efficacy**
Tobias Neuhann

SUNDAY SEPT. 8th

- 10:45 **The Glaucoma Surgical Journey:
Prioritizing Safety with
iStent *inject*[®] W**
José M^a Martínez de la Casa
- 11:15 **Interventional Glaucoma:
Beyond Understanding
to Action**
Ike Ahmed
- 13:00 **Interventional Glaucoma:
Acting Early to Save Vision**
Anselm Jünemann
- 15:00 **Less Risk:
Glaucoma Management
with iStent *inject*[®] W**
Ana Miguel

GLAUKOS

1- Gillmann K, Hombek DM. BMJ Open Opth 2024;9:e001575.doi:10.1136/bmjophth-2023-001575.

2- Samuelson, Thomas W., et al. "Quality of life in primary open-angle glaucoma and cataract: an analysis of VFQ-25 and OSDI from the iStent *inject*[®] pivotal trial." American Journal of Ophthalmology 229 (2021): 220-229.

iStent *inject*[®] W IMPORTANT SAFETY INFORMATION

INDICATION FOR USE: The iStent *inject* W, is intended to reduce intraocular pressure safely and effectively in patients diagnosed with primary open-angle glaucoma, pseudo-exfoliative glaucoma or pigmentary glaucoma. The iStent *inject* W, can deliver two (2) stents on a single pass, through a single incision. The implant is designed to stent open a passage through the trabecular meshwork to allow for an increase in the facility of outflow and a subsequent reduction in intraocular pressure. The device is safe and effective when implanted in combination with cataract surgery in those subjects who require intraocular pressure reduction and/or would benefit from glaucoma medication reduction. The device may also be implanted in patients who continue to have elevated intraocular pressure despite prior treatment with glaucoma medications and conventional glaucoma surgery. CONTRAINDICATIONS: The iStent *inject* W System is contraindicated under the following circumstances or conditions: • In eyes with primary angle closure glaucoma, or secondary angle-closure glaucoma, including neovascular glaucoma, because the device would not be expected to work in such situations. • In patients with retrolubar tumor, thyroid eye disease, Sturge-Weber Syndrome or any other type of condition that may cause elevated episcleral venous pressure. WARNINGS/ PRECAUTIONS: • For prescription use only. • This device has not been studied in patients with uveitic glaucoma. • Do not use the device if the Tyvek[®] lid has been opened or the packaging appears damaged. In such cases, the sterility of the device may be compromised. • Due to the sharpness of certain injector components (i.e. the insertion sleeve and trocar), care should be exercised to grasp the injector body. Dispose of device in a sharps container. • iStent *inject* W is MR-Conditional; see MRI Information below. • Physician training is required prior to use of the iStent *inject* W System. • Do not re-use the stent(s) or injector, as this may result in infection and/or intraocular inflammation, as well as occurrence of potential postoperative adverse events as shown below under "Potential Complications." • There are no known compatibility issues with the iStent *inject* W and other intraoperative devices. (e.g., viscoelastics) or glaucoma medications. • Unused product & packaging may be disposed of in accordance with facility procedures. Implanted medical devices and contaminated products must be disposed of as medical waste. • The surgeon should monitor the patient postoperatively for proper maintenance of intraocular pressure. If intraocular pressure is not adequately maintained after surgery, the surgeon should consider an appropriate treatment regimen to reduce intraocular pressure. • Patients should be informed that placement of the stents, without concomitant cataract surgery in phakic patients, can enhance the formation or progression of cataract. ADVERSE EVENTS: Please refer to Directions For Use for additional adverse event information. CAUTION: Please reference the Directions For Use labelling for a complete list of contraindications, warnings and adverse events.